





Director's Review of SoLID

February 10, 2021

PVDIS Theory




Jens Erler



Electroweak precision physics

- * The electroweak (EW) precision program started about 50 years ago
- * M_W , M_Z , m_t , M_H (and m_c) have all been successfully predicted before their discoveries
- * 2012 the Standard Model (SM) was completed ...
... and it is as successful as it is unsatisfactory (dark matter, naturalness, ...)
- * so far no new states discovered at the LHC, so perhaps they show up in EW physics first
- * currently some tensions in $g_{\mu-2}$, M_W , and the first row CKM matrix unitarity constraint
- * General remark: the higher the precision, the more physics issues will enter in the interpretation of precision measurements
- * this is an obstacle when looking at single observables but may be rather a feature in global analyses (across different observables and subfields of particle, nuclear and atomic physics)
- * for SoLID: mostly EW and QCD (higher twist and PDFs)

Importance of $\sin^2\theta_W$



indirect m_t


176.4 ± 1.9 GeV

(1.9 σ high)


$$\sin^2 \theta_W = \frac{g'^2}{g^2 + g'^2}$$

Freitas & JE, PDG 2020

Weak Mixing Angle and Boson Masses



$\sin^2\theta_W$ measurements



LEP & SLC:
 0.23151 ± 0.00016


Tevatron:
 0.23148 ± 0.00033

LHC:
 0.23131 ± 0.00033

average direct
 0.23148 ± 0.00013

global fit
 0.23155 ± 0.00004


Running weak mixing angle



courtesy of
Rodolfo Ferro,
see also
Ferro-Hernández, JE
arXiv:1712.09146

$$\Delta \sin^2\theta_W = \pm 0.00057$$

Beyond $\sin^2\theta_W$: discriminating new physics



- * **Z-Z' mixing:** modification of Z vector coupling
- * **oblique parameters:** STU (also need M_W and Γ_Z)
- * **new amplitudes:** off- versus on-Z pole measurements (e.g. heavy Z')
- * **dark Z:** renormalization group evolution (low versus very low energy measurements)

Standard Model Effective Field Theory (SMEFT)

- * Systematic expansion in inverse powers of new physics mass scale $\Lambda \gg M_Z$
- * no known a priori reason to stop at the level of renormalizable interactions ($D = 4$)
- * ν oscillations accounted for by 2 (12) $D = 5$ Weinberg 1979 $H^2 L^2 + \text{H.c.}$ ($\Delta L = \pm 2$) operators for 1 (3) fermion generations, counting Hermitian conjugates
- * 15 bosonic + 38 fermionic + 31 mixed = 84 (3045) independent $D = 6$ operators (Λ^{-2})
Grzadkowski et al., arXiv:1008.4884
- * 38 fermionic operators = 3 L^4 + 13 $L^2 Q^2$ + 8 $L Q^3$ ($\Delta B \neq 0$) + 14 Q^4 operators
- * 3 L^4 = $e_\nu e_\nu$ + $e_A e_\nu$ (MOLLER) + $e_A e_A$
- * 13 $L^2 Q^2$ = 7 vector and axial-vector combinations + 4 scalar + 2 tensor
- * 2 $e_\nu q_\nu$ (C_0) + 2 $e_A q_\nu$ (C_1) (APV, Qweak, P2) + 2 $e_\nu q_A$ (C_2) (SoLID) + 2 $e_A q_A$ (C_3) (e^+ @SoLID)
-1 constraint $(\bar{u}_L \gamma^\mu u_L - \bar{d}_L \gamma^\mu d_L) \bar{e}_R \gamma_\mu e_R = 0$

Electroweak physics with SoLID

$$* A_{LR}^{eDIS} \approx -\frac{9}{20\pi\alpha(Q)} \frac{Q^2}{v^2} \left[\left(\frac{2}{3}g_{AV}^{eu} - \frac{1}{3}g_{AV}^{ed} \right) + \left(\frac{2}{3}g_{VA}^{eu} - \frac{1}{3}g_{VA}^{ed} \right) \frac{1 - (1-y)^2}{1 + (1-y)^2} \right]$$

$$* A_{LR}^{eDIS} \approx 5 \times 10^{-4} (\gg A_{LR}^{MOLLER})$$


* iso-scalar (deuterium) target

* > 90% longitudinal polarization

* polarimetry: $\Delta P/P \lesssim 0.4\%$

* total systematic $\approx 0.5\%$



EW physics



SMEFT and the LHC

- * Sensitivity to new physics $D = 6$ operators through interference $(D = 6) \otimes \text{SM}$
- * Λ^{-4} effects are negligible at $Q^2 \ll M_Z^2$, i.e. in any fixed target experiment
- * Drell-Yan lepton pair production is a high-precision tool at the LHC
- * $\sin^2\theta_W$ from $A_{\text{FB}}(e, \mu)$ in a ± 30 GeV window around M_Z
- * in very high Q^2 Drell-Yan **production cross-section** data
 - $\sum_i |D = 6|^2 \geq 0$ enters at the order (Λ^{-4}) of $D = 8$ operators $(D = 8) \otimes \text{SM}$
- * in principle they constrain **all** $D = 6$ operators at $Q^2 \gg M_Z^2$
- * however, there are 993 (44807) $D = 8$ operators *Henning et al., arXiv:1512.03433*
- * effectively, they introduce an extra theory uncertainty on the $D = 6$ LHC constraints arising dominantly from total cross-sections and A_{FB} *Alte et al., arXiv:1812.07575*

Parity-violating 4-fermion electron-quark couplings




JE et al., arXiv:1401.6199

Scale exclusion from PVDIS and SoLID


New physics scale: $\frac{g}{2v^2} \rightarrow \left[\frac{g}{2v^2} + \frac{4\pi}{\Lambda^2} \right]$

$$\Lambda \gtrsim v \sqrt{\frac{2.92 \times 8\pi}{1.96 \times \Delta[2g_{AV}^{eu} - g_{AV}^{ed} + 0.84(2g_{VA}^{eu} - g_{VA}^{ed})]}}$$

$\approx 22 \text{ TeV (95 \% CL)}$




Lifting flat LHC directions



D = 8 operator

$$C_{l^2 q^2 D^2}^{(1)} \left[\frac{1}{\text{TeV}^4} \right]$$



courtesy of Frank Petriello ; see also Boughezal et al. arXiv:2004.00748


Leptophobic Z's

- * Extra Z bosons one of the most well-motivated new physics scenarios
- * very strong mass limits from the LHC, but simplified analyses allow loopholes
- * e.g., leptophobic Z's decaying into supersymmetric or dark matter particles need different search strategies *González-Alonso et al., arXiv:1211.4581*

- * $M_{Z'} \gtrsim 800$ GeV from precision data from ZZ'-mixing for the unique leptophobic Z' from E₆,
but $\theta_{ZZ'} \propto C \frac{M_Z}{M_{Z'}}$

where C can be tuned to vanish

JE et al., arXiv:0906.2435




Pérez et al., arXiv:2003.09426

Conclusions

- * ν DIS very important in establishing the SM, but the complicated physics of heavy nuclei presents bottle neck to high precision; the scepter is handed on to eDIS
- * SoLID will be perfectly synchronized with ultra-high precision PVES with P2 (Mainz) and MOLLER, APV (including isotope ratios) and precision CEvNS
- * PVDIS with SoLID precision at sub-% level
- * no convincing new physics signal at LHC yet: need to look under each rock lamppost
- * PVDIS provides such a lamppost (a concrete direction in SMEFT operator space)
- * viable models relevant to SoLID need tuning (nowadays a generic feature in NP searches); parameter space becomes fractal (each available piece of parameter space unlikely and contrived, but probably many of these)
- * but SoLID explores directions in SMEFT parameter space to which the LHC is blind

Thank You



Backup

New Physics scales Λ_{NP} (95% CL)

	precision	$\Delta\sin^2\theta_W$	Λ_{NP}		precision	$\Delta\sin^2\theta_W$	Λ_{NP}
E158	14 %	0,0013	17.0 TeV	MOLLER	2,4 %	0,00028	38 TeV
PVDIS	4,1 %	0,0043	7.8 TeV	SoLID	0,6 %	0,00057	22 TeV
Qweak	6,3 %	0,0011	27.8 TeV	P2	1,83 %	0,00033	51 TeV
				P2 ^{12}C	0,3 %	0,0007	49 TeV
APV ^{133}Cs	0,58 %	0,0019	32.3 TeV	APV ^{225}Ra	0,5 %	0,0018	34 TeV
$^{176}\text{Yb}/^{170}\text{Yb}$	0,78 %	0,052	4.3 TeV	$^{225}\text{Ra}/^{213}\text{Ra}$	0,1 %	0,0037	16 TeV